

# Approaching the molecular origins of collective dynamics in oscillating cell populations

Pankaj Mehta<sup>1</sup> and Thomas Gregor<sup>2</sup>

From flocking birds, to organ generation, to swarming bacterial colonies, biological systems often exhibit collective behaviors. Here, we review recent advances in our understanding of collective dynamics in cell populations. We argue that understanding population-level oscillations requires examining the system under consideration at three different levels of complexity: at the level of isolated cells, homogenous populations, and spatially structured populations. We discuss the experimental and theoretical challenges this poses and highlight how new experimental techniques, when combined with conceptual tools adapted from physics, may help us overcome these challenges.

## Addresses

<sup>1</sup> Department of Molecular Biology, Princeton University, Princeton, NJ 08544, USA

<sup>2</sup> Department of Physics and the Lewis-Sigler Institute for Integrative Genomics, Princeton University, Princeton, NJ 08544, USA

Corresponding author: Gregor, Thomas ([tg2@Princeton.EDU](mailto:tg2@Princeton.EDU))

**Current Opinion in Genetics & Development** 2010, **20**:574–580

This review comes from a themed issue on  
Genetics of system biology  
Edited by Jeff Hasty, Alex Hoffmann and Susan Golden

Available online 9th October 2010

0959-437X/\$ – see front matter

© 2010 Elsevier Ltd. All rights reserved.

DOI [10.1016/j.gde.2010.09.004](https://doi.org/10.1016/j.gde.2010.09.004)

## Introduction

Collective behaviors are ubiquitous in biological systems. At the molecular level, proteins often aggregate into self-organized structures such as spindles [1]. At the cellular level, unicellular organisms often form structured communities composed of many individuals [2–4]. At the level of organisms, birds and fish colonies exhibit dramatic emergent behaviors such as flocking [5–7] and schooling [8–10] (see Figure). Our understanding of collective behaviors in biological systems, however, is still in its infancy, highlighting the crucial need to study systems where the link between macroscopic behavior and the microscopic components that make up the system can be probed directly through experiments. This review focuses on one class of systems where such an approach is possible: the collective dynamics in cellular populations.

In their natural environments, cells often undertake complex collective behaviors in response to environmen-

tal and population cues [11,12]. Thus, understanding how cells behave in the wild requires characterizing not only the behavior of isolated cells but also how environmental signals combine with cell-to-cell communication (such as quorum sensing [13] and autocrine signaling [14]) to give rise to observed behaviors at the population level. Doing so requires us to examine how the cooperative behaviors of cell colonies differ from those of isolated cells and conversely, how the properties of single cells generate and explain the observed communal behavior.

The challenges inherent in this research program are summarized by Phil Anderson's famous declaration "More is Different" [15]—namely, systems composed of many interacting components will exhibit new emergent behaviors that cannot be understood simply by looking at the behavior of the individual components that make up the system. Whereas Anderson largely had in mind physical systems, biological systems pose additional challenges not encountered in physics. The collective behavior of cellular populations often require cells to integrate information from a wide variety of sources in order to perform a desired task such as cellular aggregation or cellular differentiation [16,17]. These challenges highlight the crucial need to simultaneously observe the behavior of individual cells within a population, the behavior of the population as a whole, and to measure the relevant signaling and environment induced interactions between cells.

## Oscillations in communicating cell populations

A particularly attractive system to study collective behaviors is provided by communicating cell populations that display rhythmic activities in the form of intracellular oscillations of signaling molecules or gene expression. Collective cellular oscillations play an important role in a wide variety of biological systems [18], ranging from neural systems [19] to the social amoebae *Dictyostelium discoideum*, where synchronized oscillations lead starved cells to aggregate [20–24,25••], to glycolytic and non-glycolytic oscillations in yeast populations [26–32], to oscillations in the pancreatic islets that control insulin secretion [33,34••]. Recently, even bacteria have been synthetically engineered to exhibit collective oscillations [35••]. Oscillations represent an especially tractable example of collective behavior because the link between macroscopic behavior and molecular interactions can be readily experimentally tested. Oscillations are easy to observe experimentally, can be unmasked even in noisy


data using analytical tools such as Fourier transforms, and there exist a large body of theoretical work to help interpret existing and guide new experiments. In addition, such systems are amenable to theoretical analysis using ideas from theory of dynamical systems [36–38].

Understanding collective oscillations requires disentangling behaviors at three different levels of complexity (see Figure 1). At the cellular level, it is necessary to characterize how cellular networks of genes and proteins allow single cells to respond to external signals (such as environmental cues and signaling molecules) as well as how these signals control the production of secreted molecules that are involved in cell-to-cell communication. The behavior of single cells must then be related to the behavior of cellular populations by exploring how system parameters, such as cell-density, change the collective dynamics. Finally, one must understand the spatial dynamics of these cellular oscillation processes.

The main challenge faced when studying such systems is to understand how behaviors at lower levels of complexity shape and give rise to the behaviors seen at higher levels of complexity [15].

Recently, this program has been carried out with some success in two systems, one natural and the other synthetic. Gregor *et al.* [25\*\*] used a FRET-sensor to measure internal levels of the signaling molecule cAMP [39,40]. They showed that isolated *Dictyostelium* amoeba behave like an excitable system, with individual cells capable of generating sustained oscillations in response to elevated levels of external cAMP. They then related these single-cell oscillations to the synchronized, cell-density-dependent oscillations exhibited by homogeneous cell populations and mapped out a phase diagram indicating under what conditions collective oscillations occur. The external cAMP level was identified as the control parameter that determines the oscillatory state of the system. Finally, they observed small populations of

Figure 1



Collective behaviors in biology at different levels of complexity. Top: Collective behaviors in biology exist at the molecular (mitotic spindle formation [1]), the cellular (social amoebae aggregation [4]), and the organismal (schooling fish [8]) levels. This review focuses on systems at the cellular level. Cellular organisms retain many of the interesting phenomena found in higher order organisms such as information processing and collective decision making, with the added advantage that behavior can be directly linked to processes at the molecular level. Bottom: Cellular systems can be analyzed at three different levels of complexity, at the level of isolated cells, homogenous cell populations, and spatially structured populations [77].

Understanding behavior requires systematic examination of these systems at all three levels of complexity. The main challenge faced when examining these systems is to link behavior at the single cell level to that of populations and vice versa.

*Dictyostelium* cells on agar where they showed that the first cell that randomly pulses entrains the rest of the population in rhythmic activity. The ensuing synchronized oscillations gave rise to spatial, concentric waves, with cells eventually aggregating at the wave origin, the spatial center from which the first oscillation pulse was emitted. What was notable about these series of experiments was that Gregor *et al.* were able to show that the system undergoes a collective transition from a non-oscillatory state where all cells are quiescent to a state where all cells oscillate synchronously. Furthermore, they showed that the transition does not result from specialized pacemaker cells, but is a direct consequence of the excitable nature of individual cells.

Using tools from synthetic biology, Danino *et al.* [35\*\*] engineered a genetic circuit in *E. coli* capable of generating synchronized oscillations in growing populations. One of the unique features of the system is that a population of cells, that in isolation is incapable of oscillating, exhibits collective oscillations when coupled using a quorum-sensing molecule. The genetic circuits utilized components of the naturally occurring quorum-sensing machinery in other bacterial species to induce a global coupling between cells [41,42]. Using cleverly designed microfluidic chambers (networks of micro-channels that house cells in fluid flow) that allow bacteria to grow naturally while simultaneously holding the cell-density fixed, Danino *et al.* were able to control the density of cells, and consequently the external concentration of signaling molecules, to induce synchronized oscillations in cell colonies. The experiments were then repeated in larger microfluidic chambers where spatial inhomogeneities resulted in a multitude of fascinating phenomena such as traveling waves and front propagation. What is groundbreaking about this work is that it provides proof of principle that one can engineer the properties of a system at the level of a single cell to control behavior at the level of cellular populations.

Both systems discussed above exhibit a cell-density-dependent transition to collective oscillations that has been termed ‘dynamical quorum sensing’ [43\*\*,44\*]. This phenomenon was first explicated in the context of glycolytic oscillations in yeast through a successful combination of theory and experiment [43\*\*]. Dynamical quorum sensing relies on the mutual synchronization of cells through the exchange of chemicals (metabolites in yeast, cAMP in *Dictyostelium*, quorum-sensing molecules for the engineered circuits discussed above). Since the cells themselves produce the chemicals, the external concentration of the chemicals reflects the local cell-density of the population. Collective oscillations emerge when the external concentration, or equivalently cell-density, exceeds some crucial threshold. Thus, in dynamical quorum sensing, cell-density information is encoded in the collective intracellular dynamical state of the entire population. Finally, it is worth noting that

the term ‘dynamical quorum sensing’ is used by various authors to refer to qualitatively different types of density-dependent transitions. This highlights the need for a better theoretical understanding of the qualitatively different ways that density-dependent transitions to synchronized oscillations can occur.

### Challenges in understanding collective behaviors

Understanding and manipulating collective behaviors in cellular systems poses a number of new experimental and theoretical challenges. On the experimental side, the advent of fluorescent markers has resulted in tremendous progress [45–47]. These markers include derivatives of various fluorescent proteins that can be genetically encoded and directly tag signaling proteins [48–53], as well as reporter constructs for smaller signaling molecules such as ions ( $\text{Ca}^{2+}$ ) [54,55] and nucleotides (ATP, cAMP, cGMP) [56,57]. These markers and sensors work very well inside cells where they are synthesized. However, understanding the signaling that underlies collective behavior also requires measuring signaling molecule concentrations in the extracellular space. This is particularly challenging when the individual cells are not packed together but free floating in solution. Possible techniques that may allow for the measurement of the extracellular, spatio-temporal dynamics of signaling molecules include engineering cells to artificially release sensors into the environment, as well as tagging the outside of the cell membranes with sensor molecules. For example in cortical astrocytes extracellular ATP release has been reported by both real-time imaging using bioluminescence [58] as well as using chemiluminescence with cell surface-tagged beads [59]. Extreme care has to be taken with these methods because genetically altering cells, generally, can simply give rise to collective behaviors that differ from those exhibited by wild-type cells. Alternatively, the experimenter could supply sensors externally by coating the walls of the experimental setup or by continuous flow in solution. Both require the highly controllable environments of miniature size provided by microfluidics [60–63]. Microfluidics refers to fluid flow in a network of micro-channels that houses cells and can be integrated on disposable, low-cost Lab-on-a-Chip cartridges [64,65]. Cells can survive in these environments, be easily tracked and still retain most of the natural characteristics necessary to probe collective behaviors. The ability to perform live-cell imaging while simultaneously measuring the spatio-temporal dynamics of both intracellular and extracellular signaling molecules is likely to greatly expand our understanding of collective behaviors in cellular colonies over the next decade.

Mathematical and computational modeling will also likely play an important role in expanding our understanding of cellular oscillations. Mathematical models have helped shape our current understanding of *Dictyos-*

Table 1

**Four roads to synchronized oscillations. 1. Specialized pacemaker cells such as in the heart [78]. 2. Phase locking and frequency locking of individually oscillating cells [79–82]. 3. Oscillator death in oscillators coupled with time delays [83,84]. 4. Dynamic quorum sensing such as in the social amoebae [20] and yeast [36]**

Type of transition	Below the transition	Mechanism
Pacemaker	Cells are excitable. Pacemakers do not fire.	Firing of specialized pacemaker cells trigger oscillations.
Kuramoto (phase locking)	Individual cells oscillate at their natural frequencies.	An increase in the coupling strength results in phase and/or frequency locking.
Oscillator Death	Isolated cells oscillate at a wide range of frequencies. However, when cells are coupled, individual cells do not oscillate.	A decrease in the strength of the coupling between oscillators leads to collective oscillations.
Dynamic Quorum Sensing	Cells do not oscillate but can become oscillatory in response to an external signal they themselves produce.	An increase in cell-density leads to a larger concentration of the external signaling molecules. When the concentration increases beyond the crucial concentration (i.e. bifurcation point), the cells start collectively oscillating.

*telium* by pointing out important connections with the theory of excitable systems—systems like neurons where a small change in inputs/parameters can elicit large responses such as a spike [66,67] or the stochastic release of the second messenger molecule Calcium that can give rise to sustained oscillations [68,69]. Mathematical models have also highlighted the importance of feedback loops and balancing time scales for oscillations [70] as well as aiding the design of genetic circuits capable of oscillations [35•,71,72].

Despite these considerable achievements, theory has not kept pace with the rapid experimental advances of the past decade. The need for new conceptual and theoretical approaches to collective behavior in biological systems becomes even more clear when we contrast our current level of understanding of biological systems to their physics counterparts [15,73]. The major theoretical challenge is to understand how the microscopic details of a system shape collective behaviors at larger scales. Though this theoretical program seems daunting, we can draw on inspiration from the study of collective behavior in physics. A unifying theme in the study of collective phenomenon in physics is the idea of ‘universality’—the idea that many collective properties depend only on a few ‘relevant’ microscopic details of the system under consideration [74]. The role of theory is to identify these relevant details and understand how they give rise to the observed behaviors at macroscopic scales. Recent work suggests that universality is also likely to be relevant to biological systems. For example, recent experiments demonstrate that despite its vast complexity, the yeast cell-cycle network exhibits phase locking in response to a periodic driving force, much like an idealized oscillator [75].

Another important implication of universality is that, often, there are only a few qualitatively different collective

behaviors a system can exhibit. For example, tools from dynamical systems such as bifurcation theory allow for a classification of the qualitatively different behaviors that can be exhibited by a neuron. Theoretical considerations also suggest that there are probably only a few different ways that cells can undergo a density-dependent, dynamical quorum-sensing transition to synchronized collective oscillations [36,38]. The accompanying table summarizes four common routes to synchronized oscillations seen in nature. A key challenge facing researchers studying cellular rhythms is to relate the type of dynamical quorum-sensing transition exhibited by a system to relevant microscopic details such as cell coupling. For example, Ref. [34•] utilized ideas from percolation theory to show that altering the gap–junction couplings between cells can qualitatively affect the emergence of collective calcium oscillations in the pancreatic islet (Table 1).

A final theoretical challenge is that, in contrast with physical systems, biological systems often use oscillations to perform a desired task in response to environmental and cellular signals. The classical example being the aggregation of *Dictyostelium* cells in response to starvation [24]. Thus, fully understanding biological rhythms requires integrating conceptual tools from statistical physics and dynamical systems with tools for understanding signal processing such as information theory [17,76].

## Conclusion and outlook

Collective oscillations in cellular systems represent a rich avenue of research for both biology and the physical sciences. It is now clear that the behavior of cellular populations arises from a complex interplay of components at the molecular and cellular levels. Understanding this behavior will require us to develop new theoretical and experimental tools linking the properties of single cells to those of cell populations. Ultimately, this

knowledge should allow us to control the behavior of entire cell populations simply by manipulating the properties of isolated cells.

## Acknowledgements

We thank Silvia De Monte, Jordi Garcia-Ojalvo, and Jeff Hasty for insightful comments on this review. Work was partially supported by NIH Grant K25GM086909 to PM and a Searle Scholarship to TG. We would also like to thank the Aspen Center for Physics where part of this work was completed.

## References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
  - of outstanding interest
1. Surrey T, Nedelec F, Leibler S, Karsenti E: **Physical properties determining self-organization of motors and microtubules.** *Science* 2001, **292**:1167.
  2. Mignot T, Kirby J: **Genetic circuitry controlling motility behaviors of *Myxococcus xanthus*.** *BioEssays* 2008, **30**:733-743.
  3. Bonner J: *The Social Amoebae: The Biology of Cellular Slime Molds* Princeton University Press; 2009.
  4. O'Toole G, Kaplan H, Kolter R: **Biofilm formation as microbial development.** *Annual Reviews in Microbiology* 2000, **54**:49-79.
  5. Ballerini M, Cabibbo N, Candelier R, Cavagna A, Cisbani E, Giardina I, Orlandi A, Parisi G, Procaccini A: **Empirical investigation of starling flocks: a benchmark study in collective animal behaviour.** *Animal Behaviour* 2008, **76**:201-215.
  6. Ballerini M, Cabibbo N, Candelier R, Cavagna A, Cisbani E, Giardina I, Lecomte V, Orlandi A, Parisi G, Procaccini A: **Interaction ruling animal collective behavior depends on topological rather than metric distance: evidence from a field study.** In *Proceedings of the National Academy of Sciences of the United States of America* 2008, **105**:1232.
  7. Cavagna A, Giardina I: **Large scale behaviour in animal groups.** *Behavioural Processes* 2010, **66**:.
  8. Shaw E: **Schooling fishes.** *American Scientist* 1978, **66**:166-175.
  9. Couzin I, Krause J, Franks N, Levin S: **Effective leadership and decision-making in animal groups on the move.** *Nature* 2005, **433**:513-516.
  10. Ward A, Sumpter D, Couzin I, Hart P, Krause J: **Quorum decision-making facilitates information transfer in fish shoals.** In *Proceedings of the National Academy of Sciences of the United States of America* 2008, **105**:6948.
  11. Shapiro J: **Thinking about bacterial populations as multicellular organisms.** *Annual Reviews in Microbiology* 1998, **52**:81-104.
  12. Shank E, Kolter R: **New developments in microbial interspecies signaling.** *Current Opinion in Microbiology* 2009, **12**:205-214.
  13. Waters CM, Bassler BL: **Quorum sensing: cell-to-cell communication in bacteria.** *Annual Review of Cell and Developmental Biology* 2005, **21**:319-346.
  14. Kawano S, Otsu K, Kuruma A, Shoji S, Yanagida E, Muto Y, Yoshikawa F, Hirayama Y, Mikoshiba K, Furuichi T: **ATP autocrine/paracrine signaling induces calcium oscillations and NFAT activation in human mesenchymal stem cells.** *Cell Calcium* 2006, **39**:313-324.
  15. Anderson PW: **More is different.** *Science* 1972, **177**:393-396.
  16. Bischofs IB, Hug JA, Liu AW, Wolf DM, Arkin AP: **Complexity in bacterial cell-cell communication: quorum signal integration and subpopulation signaling in the *Bacillus subtilis* phosphorelay.** In *Proceedings of the National Academy of Sciences of the United States of America* 2009, **106**:6459-6464.
  17. Mehta P, Goyal S, Long T, Bassler BL, Wingreen NS: **Information processing and signal integration in bacterial quorum sensing.** *Molecular Systems Biology* 2009, **5**:325.
  18. Murray J: *Mathematical Biology: An Introduction* Springer Verlag; 2002.
  19. Koizumi S: **Synchronization of Ca<sup>2+</sup> oscillations: involvement of ATP release in astrocytes.** *FEBS Journal* 2009, **277**:286-292.
  20. Gerisch G, Hess B: **Cyclic-AMP-controlled oscillations in suspended *Dictyostelium* cells: their relation to morphogenetic cell interactions.** In *Proceedings of the National Academy of Sciences of the United States of America* 1974, **71**:2118.
  21. Devreotes P, Potel M, MacKay S: **Quantitative analysis of cyclic AMP waves mediating aggregation in *Dictyostelium discoideum*.** *Developmental Biology* 1983, **96**:405-415.
  22. Siegert F, Weijer C: **Analysis of optical density wave propagation and cell movement in the cellular slime mould *Dictyostelium discoideum*.** *Physica D: Nonlinear Phenomena* 1991, **49**:224-232.
  23. Nagano S: **Diffusion-assisted aggregation and synchronization in *Dictyostelium discoideum*.** *Physical Review Letters* 1998, **80**:4826-4829.
  24. Sawai S, Thomason P, Cox E: **An autoregulatory circuit for long-range self-organization in *Dictyostelium* cell populations.** *Nature* 2005, **433**:323-326.
  25. Gregor T, Fujimoto K, Masaki N, Sawai S: **The onset of collective behavior in social amoebae.** *Science* 2010, **328**:1021-1025. Shows that a density-dependent transition and not specialized pacemaker cells lead populations of quorum-sensing *Dictyostelium* cells to synchronized oscillations.
  26. Hess B, Boiteux A, K $\ddot{u}$ ger J: **Cooperation of glycolytic enzymes.** *Advances in Enzyme Regulation* 1969, **7**:149-167.
  27. Dan $\ddot{o}$  S, Sørensen PG, Hynne F: **Sustained oscillations in living cells.** *Nature* 1999, **402**:320-322.
  28. Richard P: **The rhythm of yeast.** *FEMS Microbiology Reviews* 2003, **27**:547-557.
  29. Palkova Z, Vachova L: **Ammonia signaling in yeast colony formation.** *International Review of Cytology* 2003, **225**:229-272.
  30. Klevecz R, Bolen J, Forrest G, Murray D: **A genomewide oscillation in transcription gates DNA replication and cell cycle.** In *Proceedings of the National Academy of Sciences of the United States of America* 2004, **101**:1200.
  31. Tu B, McKnight S: **Metabolic cycles as an underlying basis of biological oscillations.** *Nature Reviews Molecular Cell Biology* 2006, **7**:696-701.
  32. Dan $\ddot{o}$  S, Madsen MF, Sørensen PG: **Quantitative characterization of cell synchronization in yeast.** In *Proceedings of the National Academy of Sciences of the United States of America* 2007, **104**:12732-12736.
  33. Rocheleau JV, Remedi MS, Granada B, Head WS, Koster JC, Nichols CG, Piston DW: **Critical role of gap junction coupled KATP channel activity for regulated insulin secretion.** *PLoS Biology* 2006, **4**:e26.
  34. Benninger RKP, Zhang M, Head WS, Satin LS, Piston DW: **Gap junction coupling and calcium waves in the pancreatic islet.** *Biophysical Journal* 2008, **95**:5048-5061. Uses a combination of theory and experiment to see how varying the gap-junction couplings between cells affects collective oscillations in the pancreatic islet.
  35. Danino T, Mondragón-Palomino O, Tsimring L, Hasty J: **A synchronized quorum of genetic clocks.** *Nature* 2010, **463**:326-330. Uses tools from synthetic biology to engineer bacterial cells that exhibit density-dependent collective oscillations.
  36. Strogatz S: *Nonlinear Dynamics and Chaos: With Applications to Physics, Biology, Chemistry, and Engineering* Westview Press; 2000.

37. Pikovsky A, Rosenblum M, Kurths J: *Synchronization: A Universal Concept in Nonlinear Sciences*, vol. 12 of *Cambridge Nonlinear Science Series* England: Cambridge; 2001.
38. M. Izhikevich E: **Dynamical Systems in Neuroscience: The Geometry of Excitability and Bursting** 2007:441.
39. Nikolaev VO, Bünemann M, Hein L, Hannawacker A, Lohse MJ: **Novel single chain cAMP sensors for receptor-induced signal propagation**. *The Journal of Biological Chemistry* 2004, **279**:37215-37218.
40. Bagorda A, Das S, Rericha EC, Chen D, Davidson J, Parent CA: **Real-time measurements of cAMP production in live Dictyostelium cells**. *Journal of Cell Science* 2009, **122**:3907-3914.
41. McMillen D, Kopell N, Hasty J, Collins J: **Synchronizing genetic relaxation oscillators by intercell signaling**. In *Proceedings of the National Academy of Sciences of the United States of America* 2002, **99**:679.
42. Garcia-Ojalvo J, Elowitz MB, Strogatz SH: **Modeling a synthetic multicellular clock: repressilators coupled by quorum sensing**. In *Proceedings of the National Academy of Sciences of the United States of America* 2004, **101**:10955-10960.
43. De Monte S, d'Ovidio F, Danø S, Sørensen PG: **Dynamical quorum sensing: population density encoded in cellular dynamics**. In *Proceedings of the National Academy of Sciences of the United States of America* 2007, **104**:18377-18381.  
Introduces the idea of a cell-density-dependent, dynamical quorum-sensing transition to collective oscillations in the context of glycolytic oscillations in yeast.
44. Taylor AF, Tinsley MR, Wang F, Huang Z, Showalter K: **Dynamical quorum sensing and synchronization in large populations of chemical oscillators**. *Science* 2009, **323**:614-617.  
Experimentally demonstrates the idea of the dynamical quorum sensing transition using a modified form of the classical Belousov-Zhabotinsky reaction.
45. Day R, Davidson M: **The fluorescent protein palette: tools for cellular imaging**. *Chemical Society Reviews* 2009, **38**:2887-2921.
46. Wiedenmann J, Oswald F, Nienhaus G: **Fluorescent proteins for live cell imaging: opportunities, limitations, and challenges**. *IUBMB Life* 2009, **61**:1029-1042.
47. Mavrikis M, Pourquie O, Lecuit T: **Lighting up developmental mechanisms: how fluorescence imaging heralded a new era**. *Development* 2010, **137**:373.
48. Tsien R: **The green fluorescent protein**. *Annual Review of Biochemistry* 1998, **67**:509-544.
49. Matz M, Fradkov A, Labas Y, Savitsky A, Zaraisky A, Markelov M, Lukyanov S: **Fluorescent proteins from nonbioluminescent Anthozoa species**. *Nature Biotechnology* 1999, **17**:969-973.
50. Shaner N, Campbell R, Steinbach P, Giepmans B, Palmer A, Tsien R: **Improved monomeric red, orange and yellow fluorescent proteins derived from *Discosoma* sp. red fluorescent protein**. *Nature Biotechnology* 2004, **22**:1567-1572.
51. Shcherbo D, Merzlyak E, Chepurnykh T, Fradkov A, Ermakova G, Solovieva E, Lukyanov K, Bogdanova E, Zaraisky A, Lukyanov S: **Bright far-red fluorescent protein for whole-body imaging**. *Nature Methods* 2007, **4**:741-746.
52. Deheyn D, Kubokawa K, McCarthy J, Murakami A, Porrachia M, Rouse G, Holland N: **Endogenous green fluorescent protein (GFP) in amphioxus**. *The Biological Bulletin* 2007, **213**:95.
53. Mishin A, Subach F, Yampolsky I, King W, Lukyanov K, Verkhusha V: **The first mutant of the *aequorea victoria* green fluorescent protein that forms a red chromophore**. *Biochemistry* 2008, **47**:4666-4673.
54. Brownlee C: **Cellular calcium imaging: so, what's new?** *Trends in Cell Biology* 2000, **10**:451-457.
55. Tian L, Hires S, Mao T, Huber D, Chiappe M, Chalasani S, Petreanu L, Akerboom J, McKinney S, Schreier E: **Imaging neural activity in worms, flies and mice with improved GCaMP calcium indicators**. *Nature Methods* 2009.
56. Nikolaev VO, Gambaryan S, Lohse MJ: **Fluorescent sensors for rapid monitoring of intracellular cGMP**. *Nature Methods* 2006, **3**:23-25.
57. Willoughby D, Cooper DMF: **Live-cell imaging of cAMP dynamics**. *Nature Methods* 2008, **5**:29-36.
58. Arcuino G, Lin J, Takano T, Liu C, Jiang L, Gao Q, Kang J, Nedergaard M: **Intercellular calcium signaling mediated by point-source burst release of ATP**. In *Proceedings of the National Academy of Sciences of the United States of America* 2002, **99**:9840.
59. Zhang Y, Phillips G, Li Q, Yeung E: **Imaging localized astrocyte ATP release with firefly luciferase beads attached to the cell surface**. *Analytical Chemistry* 2008, **80**:9316-9325.
60. Brody JP, Yager P, Goldstein RE, Austin RH: **Biotechnology at low Reynolds numbers**. *Biophysical Journal* 1996, **71**:3430-3441.
61. Whitesides GM: **The origins and the future of microfluidics**. *Nature* 2006, **442**:368-373.
62. Abkarian M, Faivre M, Horton R, Smistrup K, Best-Popescu CA, Stone HA: **Cellular-scale hydrodynamics**. *Biomedical Materials (Bristol, England)* 2008, **3**:034011.
63. Bennett M, Hasty J: **Microfluidic devices for measuring gene network dynamics in single cells**. *Nature Reviews Genetics* 2009, **10**:628-638.
64. Figeys D, Pinto D: **Lab-on-a-Chip: a revolution in biological and medical sciences**. *Analytical Chemistry* 2000, **72**:330-335.
65. Beebe D, Mensing G, Walker G: **Physics and applications of microfluidics in biology**. *Annual Review of Biomedical Engineering* 2003, **4**:261-286.
66. Tyson J, Keener J: **Singular perturbation-theory of traveling waves in excitable media**. *Physica D* 1988, **32**:327-361.
67. Levine H, Aranson I, Tsimring L, Truong TV: **Positive genetic feedback governs cAMP spiral wave formation in Dictyostelium**. In *Proceedings of the National Academy of Sciences of the United States of America* 1996, **93**:6382-6386.
68. Skupin A, Falcke M: **Statistical properties and information content of calcium oscillations**. *Genome Informatics* 2007, **18**:44-53.
69. Skupin A, Falcke M: **From puffs to global Ca signals: how molecular properties shape global signals**. *Chaos* 2009, **19**:037111.
70. Novak B, Tyson JJ: **Design principles of biochemical oscillators**. *Nature Reviews. Molecular Cell Biology* 2008, **9**:981-991.
71. Elowitz MB, Leibler S: **A synthetic oscillatory network of transcriptional regulators**. *Nature* 2000, **403**:335-338.
72. Süel GM, Kulkarni RP, Dworkin J, Garcia-Ojalvo J, Elowitz MB: **Tunability and noise dependence in differentiation dynamics**. *Science* 2007, **315**:1716-1719.
73. Strogatz S: **From Kuramoto to Crawford: exploring the onset of synchronization in populations of coupled oscillators**. *Physica D* 2000, **143**:1-20.
74. Kadanoff L: **Scaling and universality in statistical physics**. *Physica A* 1990, **163**:1-14.
75. Charvin G, Cross FR, Siggia ED: **Forced periodic expression of G1 cyclins phase-locks the budding yeast cell cycle**. In *Proceedings of the National Academy of Sciences of the United States of America* 2009, **106**:6632-6637.
76. Tkacik G, Callan CG, Bialek W: **Information capacity of genetic regulatory elements**. *Physical Review. E, Statistical, Nonlinear, and Soft Matter Physics* 2008, **78**:011910.
77. Basu S, Gerchman Y, Collins C, Arnold F, Weiss R: **A synthetic multicellular system for programmed pattern formation**. *Nature* 2005, **434**:1130-1134.
78. Chen P, Joung B, Shinohara T, Das M, Chen Z, Lin S: **The initiation of the heart beat**. *Circulation Journal: Official Journal of the Japanese Circulation Society* 2009.

79. Liu C, Weaver D, Strogatz S, Reppert S: **Cellular construction of a circadian clock: period determination in the suprachiasmatic nuclei.** *Cell* 1997, **91**:855-860.
80. Siapas A, Lubenov E, Wilson M: **Prefrontal phase locking to hippocampal theta oscillations.** *Neuron* 2005, **46**: 141-151.
81. To T, Henson M, Herzog E, Doyle F: **A molecular model for intercellular synchronization in the mammalian circadian clock.** *Biophysical Journal* 2007, **92**:3792-3803.
82. Ullner E, Buceta J, Diez-Noguera A, Garcia-Ojalvo J: **Noise-induced coherence in multicellular circadian clocks.** *Biophysical Journal* 2009, **96**:3573.
83. Mirolo R, Strogatz S: **Amplitude death in an array of limit-cycle oscillators.** *Journal of Statistical Physics* 1990, **60**:245-262.
84. Ullner E, Zaikin A, Volkov E, Garcia-Ojalvo J: **Multistability and clustering in a population of synthetic genetic oscillators via phase-repulsive cell-to-cell communication.** *Physical Review Letters* 2007, **99**:148103.